# HAT-P-13b,c: A TRANSITING HOT JUPITER WITH A MASSIVE OUTER COMPANION ON AN ECCENTRIC $ORBIT^{\dagger}$

G. Á. BAKOS<sup>1,2</sup>, A. W. HOWARD<sup>3</sup>, R. W. NOYES<sup>1</sup>, J. HARTMAN<sup>1</sup>, G. TORRES<sup>1</sup>, GÉZA KOVÁCS<sup>4</sup>, D. A. FISCHER<sup>5</sup>, D. W. LATHAM<sup>1</sup>, J. A. JOHNSON<sup>6</sup>, G. W. MARCY<sup>3</sup>, D. D. SASSELOV<sup>1</sup>, R. P. STEFANIK<sup>1</sup>, B. SIPŐCZ<sup>1,7</sup>, GÁBOR KOVÁCS<sup>1</sup>, G. A. ESQUERDO<sup>1</sup>, A. PÁL<sup>4,1</sup>, J. LÁZÁR<sup>8</sup>, I. PAPP<sup>8</sup>, P. SÁRI<sup>8</sup>

Draft version July 21, 2009- VERSION 06

# ABSTRACT

We report on the discovery of a planetary system with a close-in transiting hot Jupiter on a near circular orbit and a massive outer planet on a highly eccentric orbit. The inner planet, HAT-P-13b, transits the bright V=10.622 G4 dwarf star GSC 3416-00543 every  $P = 2.916260 \pm 0.000010$  days, with transit epoch  $T_c = 2454779.92979 \pm 0.00038$  (BJD) and duration  $0.1345 \pm 0.0017$  d. The outer planet, HAT-P-13c orbits the star with  $P_2 = 428.5 \pm 3.0$  days and nominal transit center (assuming zero impact parameter) of  $T_{2c} = 2454870.4 \pm 1.8$  (BJD) or time of periastron passage  $T_{2,peri} = 2454890.05 \pm 0.48$ (BJD). Transits of the outer planet have not been observed, and may not be present. The host star has a mass of  $1.22^{+0.05}_{-0.10} M_{\odot}$ , radius of  $1.56 \pm 0.08 R_{\odot}$ , effective temperature  $5638 \pm 90$  K, and is rather metal rich with  $[Fe/H] = +0.43 \pm 0.08$ . The inner planetary companion has a mass of  $0.851^{+0.029}_{-0.046} M_J$ , and radius of  $1.280 \pm 0.079 R_J$  yielding a mean density of  $0.499^{+0.103}_{-0.069}$  g cm<sup>-3</sup>. The outer companion has  $m_2 \sin i_2 = 15.2 \pm 1.0 M_J$ , and orbits on a highly eccentric orbit of  $e_2 = 0.691 \pm 0.018$ . While we have not detected significant transit timing variations of HAT-P-13b, due to gravitational and light-travel time effects, future observations will constrain the orbital inclination of HAT-P-13c, along with its mutual inclination to HAT-P-13b. The HAT-P-13 (b.c) double-planet system may prove extremely valuable for theoretical studies of the formation and dynamics of planetary systems.

Subject headings: planetary systems — stars: individual (HAT-P-13, GSC 3416-00543) techniques: spectroscopic, photometric

## 1. INTRODUCTION

Radial velocity (RV) surveys have shown that multipleplanet stellar systems are common. For example, Wright et al. (2007) concluded that the occurrence of additional planets among stars already having one known planet must be greater than 30%. Thus, one would expect that out of the  $\sim$  50 published transiting extrasolar planet (TEP) systems<sup>10</sup>, there should be a number of systems with additional companions; these companions should make their presence known through radial velocity variations of the parent stars, even if they do not themselves transit. The fact that so far no TEPs in multiple planet systems have been reported is somewhat surprising based on the statistics from RV surveys (see also Fabrycky 2009). Recently Smith et al. (2009) searched the light curves of 24 transiting planets for outer compan-

 $^{1}$  Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, gbakos@cfa.harvard.edu

NSF Fellow

<sup>3</sup> Department of Astronomy, University of California, Berkeley, CA <sup>4</sup> Konkoly Observatory, Budapest, Hungary Characteristics and Astronomy,

<sup>5</sup> Department of Physics and Astronomy, San Francisco State University, San Francisco, CA

<sup>6</sup> Institute for Astronomy, University of Hawaii, Honolulu, HI 96822; NSF Postdoctoral Fellow

<sup>7</sup> Department of Astronomy, Eötvös Loránd University, Budapest, Hungary.

Hungarian Astronomical Association, Budapest, Hungary

<sup>†</sup> Based in part on observations obtained at the W. M. Keck Observatory, which is operated by the University of California and the California Institute of Technology. Keck time has been granted by NOAO (A146Hr,A264Hr) and NASA (N128Hr,N145Hr).

 $^{10}$  www.exoplanet.eu

ions, finding no evidence for a double transiting system.

The Hungarian-made Automated Telescope Network (HATNet) survey (Bakos et al. 2002, 2004) has been a major contributor to the discovery of TEPs. Operational since 2003, it has covered approximately 10% of the Northern sky, searching for TEPs around bright stars  $(8 \leq I \leq 12.5 \text{ mag})$ . HATNet operates six wide field instruments: four at the Fred Lawrence Whipple Observatory (FLWO) in Arizona, and two on the roof of the Submillimeter Array hangar (SMA) of SAO at Hawaii. Since 2006, HATNet has announced and published 12 TEPs. A study similar to that of Smith et al. (2009) has been carried out on 9 known transiting planets from the HATNet project by Fabrycky (private communication) with a similar null result.

In this work we report on the 13th discovery of HAT-Net, the detection of the first known system with a transiting inner planet (HAT-P-13b) which also contains a second, outer planet (HAT-P-13c), as detected by the radial velocity variation of the host star. There have been examples of transiting systems where the RV variations do show a long term trend, such as HAT-P-11b (Bakos et al. 2009), but no orbital solution has vet been presented for any of these outer companions, simply because there has not been a long enough timespan to cover the period. or at least observe the long term trend changing sign. While no transits of HAT-P-13c have yet been detected, the probability that the companion actually transits the star is non-negligible if the orbits of the two planets are nearly co-planar (the pure geometric transit probability for HAT-P-13c is 1.3%, see later). The system is particularly interesting because the outer planet has both a high eccentricity and a very high mass. These properties, in turn, should induce transit timing variations (TTVs) of the inner planet of the order of 10 seconds (standard deviation, Agol et al. 2005). Such TTVs may be used to constrain the orbital parameters of the outer planet, including the inclination with respect to the line of sight and with respect to the orbital plane of the inner planet.

In § 2 of this paper we summarize the observations, including the photometric detection, and follow-up observations. § 3 describes the analysis of the data, such as the stellar parameter determination (§ 3.1), blend modeling (§ 3.2), and global modeling of the data (§ 3.3). We discuss our findings in § 4.

## 2. OBSERVATIONS

#### 2.1. Photometric detection

The transits of HAT-P-13b were detected with the HAT-5 telescope. The region around GSC 3416-00543, a field internally labeled as 136, was observed on a nightly basis between 2005 November 25 and 2006 May 20, whenever weather conditions permitted. We gathered 4021 exposures of 5 minutes at a 5.5-minute cadence. Each image contained approximately 20000 stars down to  $I \sim 14.0$ . For the brightest stars in the field we achieved a per-image photometric precision of 3.1 mmag.



FIG. 1.— The unbinned light curve of HAT-P-13 including all 4021 instrumental *I*-band measurements obtained with the HAT-5 telescope of HATNet (see text for details), and folded with the period of P = 2.9162595 days (which is the result of the fit described in § 3).

The calibration of the HATNet frames was done utilizing standard procedures. The calibrated frames were then subjected to star detection and astrometry, as described in Pál & Bakos (2006). Aperture photometry was performed on each image at the stellar centroids derived from the 2MASS catalog (Skrutskie et al. 2006) and the individual astrometrical solutions. Description of numerous details related to the reduction are also given in Pál (2009). The resulting light curves were decorrelated against trends using the External Parameter Decorrelation technique in "constant" mode (EPD, see Bakos et al. 2009) and the Trend Filtering Algorithm (TFA, see Kovács et al. 2005). The light curves were searched for periodic box-like signals using the Box Least Squares method (BLS, see Kovács et al. 2002). We detected a significant signal in the light curve of GSC 3416-00543 (also known as 2MASS 08393180+4721073;  $\alpha$  =  $08^{h}39^{m}31.82s, \ \delta = +47^{\circ}21'07.4''; \ J2000), \ with a \ depth$ of ~ 6.2 mmag, and a period of P = 2.9163 days. The dip had a relative duration (first to last contact) of  $q \approx 0.0461 \pm 0.0006$ , corresponding to a total duration of  $Pq \approx 3.228 \pm 0.040$  hours (see Fig. 1).

#### 2.2. Reconnaissance Spectroscopy

One of the important tools used in our survey for establishing whether the transit-feature in the light curve of a candidate is due to astrophysical phenomena other than a planet transiting a star is the CfA Digital Speedometer (DS; Latham 1992), mounted on the FLWO 1.5 m telescope. This yields high-resolution spectra with low signal-to-noise ratio sufficient to derive radial velocities with moderate precision (roughly  $1 \text{ km s}^{-1}$ ), and to determine the effective temperature and surface gravity of the host star. With this facility we are able to weed out certain false alarms, such as eclipsing binaries and multiple stellar systems.

We obtained 7 spectra spanning 37 days with the DS. The RV measurements of HAT-P-13 showed an rms residual of  $0.68 \text{ km s}^{-1}$ , consistent with no detectable RV variation. The spectra were single-lined, showing no evidence for more than one star in the system. Atmospheric parameters for the star, including the initial estimates of effective temperature  $T_{\text{eff}\star} = 5500 \pm 100 \text{ K}$ , surface gravity log  $g_{\star} = 4.0 \pm 0.25$  (log cgs), and projected rotational velocity  $v \sin i = 1.0 \pm 1.0 \text{ km s}^{-1}$ , were derived as described by Torres et al. (2002). The effective temperature and surface gravity correspond to a G4 dwarf (Cox 2000). The mean line-of-sight velocity of HAT-P-13 is  $+14.69 \pm 0.26 \text{ km s}^{-1}$ .

# 2.3. High resolution, high S/N spectroscopy and the search for radial velocity signal components

Given the significant transit detection by HATNet, and the encouraging DS results, we proceeded with the follow-up of this candidate by obtaining high-resolution and high S/N spectra to characterize the radial velocity variations and to determine the stellar parameters with higher precision. Using the HIRES instrument (Vogt et al. 1994) on the Keck I telescope located on Mauna Kea, Hawaii, we obtained 32 exposures with an iodine cell, plus two iodine-free templates. We used the second, higher quality template in the analysis. The observations were made between 2008 March 22 and 2009 June 5. The relative radial velocity measurements are listed in Tab. 1, and shown in Fig. 2.

Observations and reductions have been carried out in an identical way to that described in earlier HATNet discovery papers, such as Bakos et al. (2009, and references therein). References for the Keck iodine cell observations and the reduction of the radial velocities are given in Marcy & Butler (1992) and Butler et al. (1996).

Initial fits of these RVs to a single-planet Keplerian orbit were quite satisfactory but soon revealed a slight residual trend that became more significant with time. This is the reason for the observations extending over more than one year, as opposed to just a few months as necessary to confirm simpler purely sinusoidal variations seen in other transiting planets discovered by HATNet. Eventually we noticed that the residual trend reversed (Fig. 2, top), a clear sign of a coherent motion most likely due to a more distant body in the same system, possibly a massive planet. Preliminary two-planet orbital solutions provided a much improved fit (although with a weakly determined outer period due to the short duration of the observations), and more importantly, held up after additional observations. With the data so far gath-



FIG. 2.- (Top) Radial-velocity measurements from Keck for HAT-P-13, along with the best 2-planet orbital fit, shown as a function of BJD (see § 3). The center-of-mass velocity has been subtracted. (Second panel) Phased residuals after subtracting the orbital fit (also see  $\S$  3). The rms variation of the residuals is about  $3.4 \,\mathrm{m\,s^{-1}}$ , requiring a jitter of  $3.0 \,\mathrm{m\,s^{-1}}$  to be added in quadrature to the individual errors to yield a reduced  $\chi^2$  of 1.0. The errorbars in this panel have been inflated accordingly. Note the different vertical scale of the panels. (Third panel) Orbit of the outer planet after subtracting the orbital fit of the inner planet from the data. Zero phase is defined by the fictitious transit midpoint of the second planet. The error-bars (inflated by the jitter) are smaller than the size of the points. (Fourth panel) Orbit of the inner planet after subtracting the orbital fit of the outer planet. Zero-phase is defined by the transit midpoint of HAT-P-13b. The error-bars (inflated by the jitter) are smaller than the size of the points. (Fifth panel) Bisector spans (BS) for the Keck spectra including the two template spectra ( $\S$  3.2). The mean value has been subtracted. (Bottom) Relative S activity index values for the Keck spectra (see Hartman et al. 2009). Open circles in the phase-plots indicate data that appears twice due to plotting outside the [0,1] phase domain.

ered and presented in this work, a false alarm probabil-

ity (FAP) analysis for the Keplerian orbit of the second planet yielded an FAP of 0.00001.

A more thorough analysis using a two-component least squares Fourier fit (see Barning 1963, for the singlecomponent case), with one component fixed to the known frequency of the short-period inner planet, also confirmed the existence of the long period component, and indicated that the latter is due to a highly non-sinusoidal motion. A number of other low frequency peaks were eliminated as being aliases yielding worse quality fits. A three-harmonics representation of the fit yielded an RMS of  $5 \text{m s}^{-1}$ , fairly close to the value expected from the formal errors. Full modeling of this complex motion of two Keplerian orbits superposed, simultaneously with the photometry, is described in our global fit of § 3.3.

## 2.4. Photometric follow-up observations



FIG. 3.— Unbinned instrumental *i*-band transit light curves, acquired with KeplerCam at the FLWO 1.2 m telescope on seven different dates. If the first full transit is assigned  $N_{tr} = 0$  (2008 November 8/9 MST, the third event from the top), then the follow-up light curves have  $N_{tr} = -68, -1, 0, 1, 24, 35, 62$  from top to bottom. Superimposed are the best-fit transit model light curves as described in § 3. In the bottom of the figure we show the residuals from the fit. Error-bars represent the photon and background shotnoise.

We observed 7 transit events of HAT-P-13 with the KeplerCam CCD on the FLWO 1.2 m telescope between 2008 April 24 and 2009 May 8. All observations were carried out in the *i*-band, and the typical exposure time was 15 seconds. The reduction of the images, including calibration, astrometry, and photometry, was performed as described in Bakos et al. (2009).

We performed EPD and TFA against trends simultaneously with the light curve modeling (for more details, see § 3 and Bakos et al. (2009)). From the series of aper-

TABLE 1 Relative radial velocity, bisector and activity index measurements of HAT-P-13.

BJD (2,454,000+)	$\begin{array}{c} \mathrm{RV^a} \\ \mathrm{(ms^{-1})} \end{array}$	$\sigma_{ m RV}{}^{ m b}_{ m (ms^{-1})}$	$_{\rm (ms^{-1})}^{\rm BS}$	$_{\rm (ms^{-1})}^{\sigma_{\rm BS}}$	$S^{c}$	$\sigma_{ m S}$
547.89534 548.80173 548.80649 548.90849 602.73395	21.42   17.80	4.30   1.43	$1.77 \\ -1.50 \\ 0.39 \\ 1.52 \\ 12.75 \\ 22$	5.85 7.27 7.14 6.73 5.35	0.0041 0.0044 0.0045 0.0041 0.0042	$\begin{array}{c} 0.00005\\ 0.00004\\ 0.00005\\ 0.00002\\ 0.00002\\ 0.00002\\ \end{array}$
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	22.56 182.48 202.33	$1.61 \\ 1.35 \\ 2.02$	$-5.63 \\ 7.35 \\ -0.37$		$0.0040 \\ 0.0042 \\ 0.0040$	$0.00002 \\ 0.00002 \\ 0.00003$

NOTE. — This table is presented in its entirety in the electronic edition of the Astrophysical Journal. A portion is shown here for guidance regarding its form and content.

<sup>a</sup> The fitted zero-point that is on an arbitrary scale (denoted as  $\gamma_{rel}$  in Tab. 4) has been subtracted from the velocities.

<sup>b</sup> The values for  $\sigma_{\rm RV}$  do not include the jitter.

<sup>c</sup> S values are on a relative scale.

TABLE 2PHOTOMETRY FOLLOW-UP OF HAT-P-13

=

$^{\mathrm{BJD}}_{(2,454,000+)}$	$\operatorname{Mag}^{a}$	$\sigma_{ m Mag}$	Mag(orig)	Filter
54581.66038	0.00550	0.00080	0.03620	i
54581.66070	0.00763	0.00080	0.03652	i
54581.66104	0.00877	0.00080	0.03686	i
54581.66138	0.00725	0.00080	0.03720	i
54581.66171	0.00670	0.00080	0.03753	i
54581.66205	0.01004	0.00080	0.03787	i
54581.66237	0.00673	0.00080	0.03819	i
54581.66271	0.00787	0.00080	0.03853	i

NOTE. — This table is presented in its entirety in the electronic edition of the Astrophysical Journal. A portion is shown here for guidance regarding its form and content.

<sup>a</sup> The out-of-transit level has been subtracted. These magnitudes have been subjected to the EPD and TFA procedures (in "ELTG" mode), carried out simultaneously with the transit fit.

tures, for each night, we chose the one yielding the smallest residual rms for producing the final light curve. These are shown in the upper plot of Fig. 3, superimposed with our best fit transit light curve models (see also  $\S$  3).

## 3. ANALYSIS

## 3.1. Properties of the parent star

We derived the initial stellar atmospheric parameters by using one of the template spectra obtained with the Keck/HIRES instrument. We used the SME package of Valenti & Piskunov (1996) along with the atomic-line database of Valenti & Fischer (2005), which yielded the following *initial* values and uncertainties (which we have conservatively increased, to include our estimates of the systematic errors): effective temperature  $T_{\text{eff}*} = 5760 \pm$ 90 K, stellar surface gravity  $\log g_* = 4.28 \pm 0.10$  (cgs), metallicity [Fe/H] =  $+0.50 \pm 0.08$  dex, and projected rotational velocity  $v \sin i = 1.9 \pm 0.5$  km s<sup>-1</sup>. Further analysis of the spectra has shown that the host star is chromospherically quiet with  $\log R'_{HK} = -5.10$  and S = 0.14(on an absolute scale).

To determine the stellar properties via a set of isochrones, we used three parameters: the stellar effective temperature, the metallicity, and the normalized semi-major axis  $a/R_{\star}$  (or related mean stellar density  $\rho_{\star}$ ). We note that another possible parameter in place of  $a/R_{\star}$  would be the stellar surface gravity, but in the case of planetary transits the  $a/R_{\star}$  quantity typically imposes a stronger constraint on the stellar models (Sozzetti et al. 2007). (The validity of this assumption, namely that the adequate physical model describing our data is a planetary transit, as opposed to e.g. a blend, is shown later in § 3.2.) Using the values of  $T_{\text{eff}\star}$ , [Fe/H], and  $\log q_{\star}$  from the SME analysis, and corresponding quadratic limb darkening coefficients by Claret (2004) (all listed in Tab. 3), we performed a global modeling of the data (§ 3.3), yielding a full Monte-Carlo distribu-tion of  $a/R_{\star}$ . This was complemented by a Monte-Carlo distribution of  $T_{\text{eff}\star}$  and [Fe/H], obtained by assuming Gaussian uncertainties based on the 1- $\sigma$  error-bars of the initial SME analysis. For each combination within the large (~ 20000) set of  $a/R_{\star}$ ,  $T_{\rm eff\star}$ , [Fe/H] values, we searched the stellar isochrones of the Yi et al. (2001)models for the best fit stellar model parameters  $(M_{\star}, R_{\star})$ age,  $\log g_{\star}$ , etc.). We derived the mean values and uncertainties of the physical parameters based on their a*posteriori* distribution (see e.g. Pál et al. 2008).

We then repeated the SME analysis by fixing the stellar surface gravity to the refined value of  $\log q_{\star} =$  $4.13 \pm 0.04$  based on the isochrone search, and only adjusting  $T_{\text{eff}\star}$ , [Fe/H] and  $v \sin i$ . This second iteration yielded  $T_{\text{eff}\star} = 5638 \pm 90 \text{ K}$ ,  $[\text{Fe}/\text{H}] = +0.43 \pm 0.08$  and  $v \sin i = 4.1 \pm 0.5 \,\mathrm{km \, s^{-1}}$ . We adopted the above values as the final atmospheric parameters for this star. We then also repeated the isochrone search for stellar parameters, yielding  $M_{\star}=1.215^{+0.051}_{-0.099} M_{\odot}$ ,  $R_{\star}=1.557 \pm 0.081 R_{\odot}$  and  $L_{\star}=2.19 \pm 0.31 L_{\odot}$ . These are summarized in Tab. 3, along with other stellar properties. Model isochrones from Yi et al. (2001) for metallicity [Fe/H] = +0.43 are plotted in Fig. 4, with the final choice of effective temperature  $T_{\rm eff\star}$  and  $a/R_{\star}$  marked, and encircled by the 1- $\sigma$  and 2- $\sigma$  confidence ellipsoids. The second SME iteration at fixed stellar surface gravity (as determined from  $a/R_{\star}$ ) changed the metallicity and stellar temperature in such a way that the new  $(T_{\mathrm{eff}\star}, a/R_{\star})$  values now fall in a more complex region of the isochrones, as compared to the initial SME values, allowing for multiple solutions

(see Fig. 4, where the original SME values are marked with a triangle).<sup>11</sup> The distribution of stellar age becomes bimodal with the dominant peak in the histogram (not shown) being at 5.0 Gyr, and a smaller peak (by a factor of 5) at 7.3 Gyr. This corresponds to a slightly bimodal mass distribution with the dominant peak at  $\sim 1.23 M_{\odot}$ , and much smaller peak around  $\sim 1.13 M_{\odot}$ . The asymmetric error-bars given in Tab. 3 for the mass and age account for the double-peaked distribution.



FIG. 4.— Stellar isochrones from Yi et al. (2001) for metallicity [Fe/H]=+0.43 and ages between 3.6 and 8.0 Gyr in steps of 0.4 Gyr. The final choice of  $T_{\rm eff\star}$  and  $a/R_{\star}$  are marked and encircled by the 1- $\sigma$  and 2- $\sigma$  confidence ellipsoids. The open triangle denotes the  $T_{\rm eff\star}, a/R_{\star}$  point found during the first SME iteration, before refining the stellar surface gravity.

The stellar evolution modeling also yields the absolute magnitudes and colors in various photometric passbands. We used the apparent magnitudes from the 2MASS catalogue (Skrutskie et al. 2006) to determine the distance of the system, after conversion to the ESO system of the models. The reported magnitudes for this star are  $J_{2MASS} = 9.328 \pm 0.018, H_{2MASS} = 9.058 \pm 0.017$ and  $K_{2MASS} = 8.975 \pm 0.017$ , which we transformed to  $J_{\rm ESO} = 9.396 \pm 0.022, H_{\rm ESO} = 9.070 \pm 0.021$  and  $K_{\rm ESO} =$  $9.018 \pm 0.018$ , following Carpenter (see 2001). These yield a color of  $(J - K) = 0.378 \pm 0.028$ , fully consistent with the expected, isochrone-based  $(J-K)_{iso} = 0.374 \pm 0.015$ . We thus relied on the 2MASS K apparent magnitude and the  $M_{\rm K} = 2.36 \pm 0.12$  absolute magnitude derived from the above-mentioned modeling to determine the distance:  $214 \pm 12 \,\mathrm{pc}$ , assuming no reddening. The K-band was chosen because it is the longest wavelength band-pass with the smallest expected discrepancies due to molecular lines in the spectrum of the star.

#### 3.2. Excluding blend scenarios

Following Torres et al. (2007), we explored the possibility that the measured radial velocities are not due to the (multiple) planet-induced orbital motion of the star, but are instead caused by distortions in the spectral line profiles. This could be due to contamination from a nearby unresolved eclipsing binary, in this case presumably with a second companion producing the RV signal corresponding to HAT-P-13c. A bisector analysis based on the Keck spectra was done as described in §3 of Hartman et al. (2009).

 TABLE 3

 Stellar parameters for HAT-P-13

Parameter	Value	Source
$T_{\rm eff\star}$ (K)	$5638\pm90$	SME <sup>a</sup>
[Fe/H] (dex)	$+0.43\pm0.08$	SME
$v \sin i  (\mathrm{km  s^{-1}})$	$4.1 \pm 0.5$	SME
$v_{\rm mac}  ({\rm km  s^{-1}})$ .	$3.78\pm0.0$	SME
$v_{\rm mic}  ({\rm km  s^{-1}})$ .	$0.85\pm0.0$	SME
$\gamma_{\rm RV}  ({\rm km  s^{-1}})$ .	$+14.69\pm0.26$	DS
$\gamma_{1(i)}$	0.3060	SME+Claret
$\gamma_{2(i)}$	0.3229	SME+Claret
$M_{\star}(M_{\odot})\dots$	$1.22^{+0.05}_{-0.10}$	$YY + a/R_{\star} + SME^{c}$
$R_{\star} (R_{\odot}) \dots \dots$	$1.56 \pm 0.08$	$YY + a/R_{\star} + SME$
$\log g_{\star} \ (\mathrm{cgs}) \ldots$	$4.13\pm0.04$	$YY+a/R_{\star}+SME$
$L_{\star} (L_{\odot}) \dots \dots$	$2.19 \pm 0.31$	$YY+a/R_{\star}+SME$
$V \pmod{\max}$	10.622	TASS <sup>d</sup>
$B-V \pmod{1}$ .	$0.73 \pm 0.03$	$YY+a/R_{\star}+SME$
$M_V \pmod{\max} \ldots$	$3.99 \pm 0.17$	$YY+a/R_{\star}+SME$
K (mag, ESO)	$8.999 \pm 0.017$	2MASS <sup>e</sup>
$M_K$ (mag,ESO)	$2.36 \pm 0.12$	$YY+a/R_{\star}+SME$
Age $(Gyr) \dots$	$5.0^{+2.5}_{-0.8}$	$YY+a/R_{\star}+SME$
Distance $(pc)$ .	$214 \pm 12$	$YY + a/R_{\star} + SME$

<sup>a</sup> SME = "Spectroscopy Made Easy" package for analysis of high-resolution spectra Valenti & Piskunov (1996). These parameters depend primarily on SME, with a small dependence on the iterative analysis incorporating the isochrone search and global modeling of the data, as described in the text.

 $^{\rm b}$  SME+Claret = Based on SME analysis and tables from Claret (2004).

<sup>c</sup> YY+ $a/R_{\star}$ +SME = Yi et al. (2001) isochrones,  $a/R_{\star}$  luminosity indicator, and SME results.

<sup>d</sup> Based on the TASS catalogue (Droege et al. 2006).

<sup>e</sup>Based on the transformations from Carpenter (2001).

We detect no significant variation in the bisector spans (see Fig. 2, fifth panel). The correlation between the radial velocity and the bisector variations is also insignificant. Therefore, we conclude that the velocity variations of the host star are real, and can be interpreted as being due to a close-in planet, with the added complication from an outer object that we account for in the modeling that follows. Because of the negligible bisector variations that show no correlation with the radial velocities, we found no need to perform detailed blend modeling of hierarchical triple (quadruple) scenarios, such as that done for the case of HAT-P-12b (Hartman et al. 2009).

#### 3.3. Global modeling of the data

This section presents a simultaneous fitting of the HATNet photometry, the follow-up light curves, and the RV observations, which we refer to as "global" modeling. It incorporates not only a physical model of the system, but also a description of systematic (instrumental) variations.

Our model for the follow-up light curves used analytic formulae based on Mandel & Agol (2002) for the eclipse of a star by a planet, where the stellar flux is described by quadratic limb-darkening. The limb darkening coefficients were taken from the tables by Claret (2004) for the *i*-band, corresponding to the stellar properties determined from the SME analysis (§ 3.1). The transit shape was parametrized by the normalized planetary radius  $p \equiv R_p/R_{\star}$ , the square of the impact parameter  $b^2$ , and the reciprocal of the half duration of the transit  $\zeta/R_{\star}$ . We chose these parameters because of their

<sup>&</sup>lt;sup>11</sup> The reason for the intersecting isochrones is the "kink" on the evolutionary tracks for  $M_{\star} \gtrsim 1.0 M_{\odot}$  stars evolving off the main sequence.

simple geometric meanings and the fact that they show negligible correlations (see e.g. Carter et al. 2008).

Our model for the HATNet data was the simplified "P1P3" version of the Mandel & Agol (2002) analytic functions (an expansion by Legendre polynomials), for the reasons described in Bakos et al. (2009). The depth of the HATNet transits was also adjusted in the fit (the depth was  $B_{inst}$  "blending factor" times that of the follow-up data) to allow for possible contamination by nearby stars in the under-sampled images of HATNet.

As indicated earlier, initial modeling of the RV observations showed deviations from a Keplerian fit highly suggestive of a second body in the system with a much longer period than the transiting planet. Thus, in our global modeling, the RV curve was parametrized by the combination of an eccentric Keplerian orbit for the inner planet with semi-amplitude K, and Lagrangian orbital elements  $(k, h) = e \times (\cos \omega, \sin \omega)$ , plus an eccentric Keplerian orbit for the outer object with  $K_2$ ,  $k_2$  and  $h_2$ , and a systemic RV zero-point  $\gamma$ . Throughout this paper the subscripts "1" and "2" will refer to HAT-P-13b and HAT-P-13c, respectively. If the subscript is omitted, we refer to HAT-P-13b.

In the past, for single transiting planet scenarios we have assumed strict periodicity in the individual transit times (Hartman et al. 2009, and references therein), even when a drift in the RV measurements indicated an outer companion (HAT-P-11b Bakos et al. 2009). Since the expected transit timing variations (TTV) for these planets were negligible compared to the error-bars of the transit center measurements, the strict periodicity was a reasonable hypothesis. Those data were characterized by two transit centers ( $T_A$  and  $T_B$ ), and all intermediate transits were interpolated using these two epochs and their corresponding  $N_{tr}$  transit numbers. The model for the RV data component also implicitly contained its ephemeris information through  $T_A$  and  $T_B$ , and thus was coupled with the photometry data in the time domain.

For HAT-P-13b, however, the disturbing force by the outer planet HAT-P-13c is expected to be not insignificant, because the RV semi-amplitude of the host star  $(\sim 0.5 \,\mathrm{km \, s^{-1}})$  indicates that HAT-P-13c is massive, and the relatively short period and eccentric orbit (see later) indicate that it moves in relatively close to HAT-P-13b. Thus, the assumption of strict periodicity for HAT-P-13b is not precisely correct. While we performed many variations of the global modeling, including assuming strict periodicity in the entire data-set, in our finally adopted physical model we assume strict periodicity only for the HATNet data, where the timing error on individual transits can be  $\sim$  1000 seconds. In this final model we allow for departure from strict periodicity for the individual transit times for the seven follow-up photometry observations. In practice, we assigned the transit number  $N_{tr} = 0$  to the first complete high quality followup light curve gathered on 2008 November 8/9 (MST). The HATNet data-set was characterized by  $T_{c,-370}$  and  $T_{c,-309}$ , covering all transit events observed by HATNet. The transit follow-up observations were characterized by their respective times of transit center:  $T_{c,-68}$ ,  $T_{c,-1}$ ,  $T_{c,0}, T_{c,1}, T_{c,24}, T_{c,35}, T_{c,62}$ . Initial estimates of the  $T_{c,i}$ values yielded an initial epoch  $T_c$  and period P by linear fitting weighted by the respective error-bars of the transit centers. The model for the RV data component

of the inner planet contained the ephemeris information through the  $T_c$  and P variables, i.e. it was coupled with the transit data. The global modeling was done in an iterative way. After an initial fit to the transit centers (and other parameters; see later),  $T_c$  and P were refined, and the fit was repeated.

The time dependence of the RV of the outer planet was described via its hypothetical transit time  $T_{2c}$  (as if its orbit were edge on), and its period  $P_2$ . The time of periastron passage  $T_{2peri}$  can be equivalently used in place of time of conjunction  $T_{2c}$ .

Altogether, the 21 parameters describing the physical model were  $T_{c,-370}$ ,  $T_{c,-309}$  (HATNet),  $T_{c,-68}$ ,  $T_{c,-1}$ ,  $T_{c,0}$ ,  $T_{c,1}$ ,  $T_{c,24}$ ,  $T_{c,35}$ ,  $T_{c,62}$  (FLWO 1.2 m),  $R_p/R_{\star}$ ,  $b^2$ ,  $\zeta/R_{\star}$ , K,  $\gamma$ ,  $k = e \cos \omega$ ,  $h = e \sin \omega$ ,  $K_2$ ,  $k_2$ ,  $h_2$ ,  $P_2$  and  $T_{2c}$ . Two additional auxiliary parameters were the instrumental blend factor  $B_{inst}$  of HATNet, and the HATNet out-of-transit magnitude,  $M_{0,\text{HATNet}}$ .



FIG. 5.— Part of the global modeling described in § 3 are corrections for systematic variations of the light curves via simultaneous fitting with the physical model of the transit. The figure shows the resulting EPD- and TFA-corrected light curves for all 7 follow-up events (small gray points), and the merged and binned light curve (bin-size 0.005 days).

We extended our physical model with an instrumental model that describes the systematic (non-physical) variations (such as trends) of the follow-up data. This was done in a similar fashion to the analysis presented in Bakos et al. (2009). The HATNet photometry had already been EPD- and TFA-corrected before the global modeling, so we only modeled systematic effects in the follow-up light curves. We chose the "ELTG" method, i.e. EPD was performed in "local" mode with EPD coefficients defined for each night, whereas TFA was performed in "global mode", with the same coefficients describing the optimal weights for the selected template stars in the field. The five EPD parameters include the hour angle (characterizing a monotonic trend that changes linearly over a night), the square of the hour angle, and three stellar profile parameters (equivalent to FWHM, elongation, and position angle). The exact functional form of the above parameters contained 6 coefficients, including the auxiliary out-of-transit magnitude of the individual events. The EPD parameters were independent for all 7 nights, implying 42 additional coefficients in the global fit. For the global TFA analysis we chose 19 template stars that had good quality measurements for all nights and on all frames, implying 19 additional parameters in the fit. Thus, the total number

of fitted parameters was 21 (physical model) + 2 (auxiliary) + 42 (local EPD) + 19 (TFA) = 84.

The joint fit was performed as described in Bakos et al. (2009), by minimizing  $\chi^2$  in the parameter space using a hybrid algorithm, combining the downhill simplex method (AMOEBA, see Press et al. 1992) with the classical linear least-squares algorithm. We exploited the fact that the partial derivatives of the model functions were derived by Pál (2008). Uncertainties in the parameters were derived using the Markov Chain Monte-Carlo method (MCMC, see Ford 2006). Since the eccentricity of the inner system appeared as marginally significant  $(k = -0.016 \pm 0.008, h = 0.008 \pm 0.012, \text{ im}$ plying  $e = 0.021 \pm 0.009$ ), and also because the physical model dictates that in the presence of a massive outer companion, the inner planet could maintain a non-zero eccentricity, we did not fix k and h to zero. The best fit results for the relevant physical parameters are summarized in Tab. 4 and Tab. 5. The individual transit centers for the FLWO 1.2 m data are given in  $\S$  4.2. Other parameters fitted but not listed in the table were:  $T_{c,-370} = 2453700.9182 \pm 0.0068$  (BJD),  $T_{c,-309} =$ 2453878.8216  $\pm$  0.0093 (BJD),  $B_{inst} = 0.82 \pm 0.06$ , and  $M_{0,\text{HATNet}} = 9.7966 \pm 0.0001$  (*I*-band). The fit to the HATNet photometry data was shown earlier in Fig. 1, the orbital fit to the RV data is shown in Fig. 2, and the fit to the FLWO 1.2 m data is exhibited in Fig. 3. The low rms of  $3.4 \,\mathrm{m \, s^{-1}}$  around the orbital fit is due in part to the use of an iodine free Keck/HIRES template that was acquired with higher spectral resolution and higher S/N than usual. Note that the low rms implies the absence of any additional interior planets other than HAT-P-13b, consistent with expectations given that the massive and eccentric outer planet HAT-P-13c dynamically forbids such planets. The stellar jitter required to reconcile the reduced  $\chi^2$  with 1.0 for the RV data is  $3.0 \,\mathrm{m \, s^{-1}}$ . The low jitter is also consistent with HAT-P-13 being a chromospherically quiet star (based on  $\log R'_{HK}$  and S).

The planetary parameters and their uncertainties can be derived by the direct combination of the *a posteriori* distributions of the light curve, radial velocity and stellar parameters. We find that the mass of the inner planet is  $M_p = 0.851^{+0.029}_{-0.046} M_J$ , the radius is  $R_p = 1.280 \pm$  $0.079 R_J$ , and its density is  $\rho_p = 0.499^{+0.103}_{-0.069} \text{ g cm}^{-3}$ . The final planetary parameters are summarized at the bottom of Table 4. The simultaneous EPD- and TFA-corrected light curve of all photometry follow-up events is shown in Fig. 5.

The outer planet, HAT-P-13c, appears to be very massive, with  $m_2 \sin i_2 = 15.2 \pm 1.0 M_J$ , and orbits the star in a highly eccentric orbit with  $e_2 = 0.691 \pm 0.018$ . The orientation of the orbit ( $\omega_2 = 176.7 \pm 0.5^\circ$ ) is such that our line of sight is almost along the minor axis, coincidentally as in the HAT-P-2b system (Bakos et al. 2007). We note that the periastron passage of HAT-P-13c has not been well monitored, and thus the RV fit of the orbit suffers from a strong correlation between the quantities  $K_2$ ,  $e_2$ and  $\gamma$ , leading to correlated  $m_2 \sin i$  and  $e_2$  (Fig. 6).

## 4. DISCUSSION



FIG. 6.— The orbital parameters, and as a consequence, the mass  $(m_2 \sin i)$  and eccentricity  $e_2$  of the outer planet are strongly correlated (see § 3). Gray dots represent the results of the MCMC runs (20,000 trials). The final solution is marked with an open circle. The bimodal distribution in  $m_2 \sin i$  is due to the similar distribution of the stellar mass (see § 3.1).

We present the discovery of HAT-P-13, the first detected multi-planet system with a transiting planet. The inner transiting planet, HAT-P-13b, is an inflated "hot Jupiter" in a nearly circular orbit. The outer planet, HAT-P-13c, is both extremely massive and highly eccentric, and orbits in a P > 1 yr orbit . With an iron abundance of  $[Fe/H] = +0.43 \pm 0.08$ , the host star is also remarkable. As we describe below, this extraordinary system is a rich dynamical laboratory for TTV effects, the first to have an accurate clock (HAT-P-13b) and a known perturbing force (HAT-P-13c).

# 4.1. The planet HAT-P-13b

The only other planet known with a host-star as metal rich as HAT-P-13 ([Fe/H]=+0.43  $\pm$  0.08) is XO-2b ([Fe/H] = 0.45, Burke et al. 2007). XO-2b, however, has much smaller mass (0.56  $M_{\rm J}$ ) and smaller radius (0.98  $R_{\rm J}$ ) than HAT-P-13b (0.85  $M_{\rm J}$ , 1.28  $R_{\rm J}$ ). Other planets similar to HAT-P-13b include HAT-P-9b ( $M_p =$  0.78  $M_{\rm J}, R_p = 1.4 R_{\rm J}$ , Shporer et al. 2008), and XO-1b ( $M_p = 0.92 M_{\rm J}, R_p = 1.21 R_{\rm J}$ , McCullough et al. 2006).

When compared to theoretical models, HAT-P-13b is a clearly inflated planet. The Baraffe et al. (2008) models with solar insolation (at  $0.045 \,\mathrm{AU}$ ) are consistent with the observed mass and radius of HAT-P-13b either with overall metal content Z=0.02 and a very young age of 0.05-0.1 Gyr, or with metal content Z=0.1 and age 0.01-0.05 Gyr.<sup>12</sup> Given the fact, however, that the host star is metal rich, and fairly old  $(5.0^{+2.5}_{-0.8} \text{ Gyr})$ , it is unlikely that HAT-P-13b is newly formed (50 Myr) and very metal poor. Comparison with Fortney et al. (2007) leads to similar conclusions. Using these models, HAT-P-13b is broadly consistent only with a 300 Myr planet at 0.02 AU solar distance and core-mass up to  $25 M_{\oplus}$ , or a 1 Gyr core-less (pure H/He) planet. If HAT-P-13b has a significant heavy rocky core, consistent with expectation given the high metallicity of HAT-P-13, it must somehow be inflated beyond models calculated for such a planet with age and insolation suggested by the data. Numerous explanations have been brought up in the past to explain the inflated radii of certain extrasolar plan-

 $<sup>^{12}</sup>$  The equivalent semi-major axis, at which HAT-P-13b would receive the same amount of insolation when orbiting our Sun, is  $a_{equiv}=0.0286^{+0.0017}_{-0.0014}$  AU.

2

8

TABLE 4 Orbital and planetary parameters for HAT-P-13b  $\,$ 

Parameter	Value			
Light curve parameters, HAT-P-13b				
P (days)	$2.916260 \pm 0.000010$			
$T_c$ (BJD)	$2454779.92979 \pm 0.00038$			
$T_{14}$ (days) <sup>a</sup>	$0.1345 \pm 0.0017$			
$T_{12} = T_{34} $ (days) <sup>a</sup>	$0.0180 \pm 0.0018$			
$a/R_{\star}$	$5.84^{+0.28}_{-0.23}$			
$\zeta/R_{\star}$	$17.07 \pm 0.16$			
$\hat{R}_p/R_\star$	$0.0844 \pm 0.0013$			
$b^{2}$	$0.447^{+0.044}_{-0.056}$			
$b = a \cos i / B$	$0.668^{+0.032}$			
i (deg)	$83.4 \pm 0.6$			
$T \rightarrow (days)$	$2454780.64 \pm 0.42$			
<i>iperi</i> (days)	2404100.04 ± 0.42			
RV parameters, as induced by HA	AT-P-13b			
$K ({\rm ms^{-1}})$	$106.1 \pm 1.4$			
<i>k</i>	$-0.016 \pm 0.008$			
h	$0.008\pm0.012$			
e	$0.021 \pm 0.009$			
$\omega$	$181 \pm 46^{\circ}$			
Other RV parameters				
$\gamma_{rel} (\mathrm{ms^{-1}}) \ldots$	$-51.3 \pm 3.8$			
Jitter $(m s^{-1})$	3.0			
Secondary eclipse parameters for HAT-P-13b				
$T_s$ (BJD)	$2454781.359 \pm 0.014$			
$T_{s,14}$	$0.1345 \pm 0.0069$			
$T_{s,12}$	$0.0180 \pm 0.0018$			
Planetary parameters for HAT-P-	13b			
$M_n(M_1)$	$0.851^{+0.029}_{-0.046}$			
$R_{n}^{P}(R_{1})$	$1.280 \pm 0.079$			
$C(M_{-}, R_{-})^{\rm b}$	0.56			
c(mp, np)	$0.400^{\pm 0.103}$			
$p_p$ (g cm )	$0.499_{-0.069}$			
<i>a</i> (AU)	$0.0426_{-0.0012}$			
$\log g_p (\text{cgs}) \dots \dots \dots \dots \dots$	$3.11 \pm 0.05$			
$T_{eq}$ (K)	$1649 \pm 45$			
$\Theta$	$0.046 \pm 0.003$			
$F_{per}$ (erg s <sup>-1</sup> cm <sup>-2</sup> ) "	$(1.74 \pm 1.9) \times 10^{+13}$			
$F_{ap} (\text{erg s}^{-1} \text{ cm}^{-2}) \in \dots \dots$	$1.6 \cdot 10^{\pm 13} \pm 1.74 \cdot 10^{\pm 12}$			
$\langle F \rangle$ (erg s <sup>-1</sup> cm <sup>-2</sup> )	$1.67 \cdot 10^{+13} \pm 1.81 \cdot 10^{+12}$			

<sup>a</sup>  $T_{14}$ : total transit duration, time between first to last contact;  $T_{12} = T_{34}$ : ingress/egress time, time between first and second, or third and fourth contact.

 $^{\rm b}$  Correlation coefficient between the planetary mass  $M_p$  and radius  $R_p.$ 

<sup>c</sup> The Safronov number is given by  $\Theta = \frac{1}{2} (V_{\rm esc}/V_{\rm orb})^2 = (a/R_p)(M_p/M_{\star})$  (see Hansen & Barman 2007).

<sup>d</sup> Incoming flux per unit surface area.

ets (Miller, Fortney, & Jackson 2009; Fabrycky, Johnson, & Goodman 2007, and references therein). One among these has been the tidal heating due to non-zero eccentricity (Bodenheimer, Lin, & Mardling 2001). No perturbing companion has been found for the inflated planets HD 209458b and HAT-P-1b (for an overview, see Mardling 2007). The HAT-P-13 system may be the first, where the inflated radius can be explained by the non-zero eccentricity of HAT-P-13b, excited by the outer HAT-P-13c planet orbiting on a highly eccentric orbit. We note that while the eccentricity of HAT-P-13b is non-zero at  $2-\sigma$ ), its pericenter is aligned (to within  $4^{\circ} \pm 40^{\circ}$ ) with that of the outer planet HAT-P-13c. Additional RV measurements and/or space-based timing of a secondary eclipse

 TABLE 5

 Orbital and planetary parameters for HAT-P-13c

Parameter	Value			
RV parameters, as induced by HAT-P-13c				
$P_2$ (days)	$428.5 \pm 3.0$			
$\bar{T_{2c}^{a}}$ (BJD)	$2454870.4 \pm 1.8$			
$K_2 (m s^{-1})$	$502 \pm 37$			
k <sub>2</sub>	$-0.690^{+0.020}_{-0.016}$			
$h_2$	$0.039 \pm 0.005$			
<i>e</i> <sub>2</sub>	$0.691 \pm 0.018$			
$\omega_2$	$176.7 \pm 0.5^{\circ}$			
$\overline{T_{2,peri}}$ (days)	$2454890.05 \pm 0.48$			
Fictitious light curve parameters, HAT-P-13c <sup>b</sup>				
$T_{2,14}^{\rm c}$ (days)	$1.283 \pm 0.059$			
$T_{2,12} = T_{34}$ (days)	$0.0734 \pm 0.0025$			
Fictitious secondary eclipse parameters for HAT-P-13c <sup>a</sup>				
$T_{2s}$ (BJD)	$2454912.7 \pm 1.8$			
$T_{2s,14}$ (days)	$1.186 \pm 0.056$			
$T_{2s,12}$ (days)	$0.0734 \pm 0.0025$			
Planetary parameters for HAT-P-13c				
$m_2 \sin i_2 (M_{\rm J}) \ldots \ldots$	$15.2 \pm 1.0$			
a2 (AU)	$1.186^{+0.018}_{-0.022}$			
$T_{2} \sim (K)$	$339 \pm 9$			
$F_{2,req}$ (rrg s <sup>-1</sup> cm <sup>-2</sup> ) d	$2.24 \cdot 10^{+11} + 3.48 \cdot 10^{+10}$			
$F_2$ , per (erg s <sup>-1</sup> cm <sup>-2</sup> ) d	$753 \cdot 10^{+09} + 8.48 \cdot 10^{+08}$			
$/F_{\rm c}$ (erg s <sup>-1</sup> cm <sup>-2</sup> ) d	$2.07 10^{+10} \pm 2.98 10^{+09}$			
\1.2/ (erg.s cill )	2.31 10 ± 3.28 10			

 $^{\rm a}$   $T_{2c}$  would be the center of transit of HAT-P-13c, if its (unknown) inclination was 90°.

 $^{\rm b}$  Transits of HAT-P-13c have not been observed. The values are for guidance only, and assume zero impact parameter.

<sup>c</sup>  $T_{14}$ : total transit duration, time between first to last contact, assuming zero impact parameter.  $T_{12} = T_{34}$ : ingress/egress time, time between first and second, or third and fourth contact. Note that these values are fictitious, and transits of HAT-P-13c have not been observed.

 $^{\rm d}$  Incoming flux per unit surface area in periastron, apastron, and averaged over the orbit.

are necessary for a more accurate determination of the orbit.

#### 4.2. Transit timing variations

It has long been known that multiple planets in the same planetary system perturb each other, and this may lead to detectable variations in the transit time of the transiting planet(s). Transit timing variations have been analytically described by Holman & Murray (2005) and Agol et al. (2005), and have been suggested as one of the most effective ways of detecting small mass perturbers of transiting planets. This has motivated extensive followup of known TEPs e.g. by the Transit Light Curve (TLC) project (Holman et al. 2006; Winn et al. 2007), looking for companions of TEPs. However, for HAT-P-13b there is observational (spectroscopic) evidence for an outer component (HAT-P-13c), and thus transit timing variations of HAT-P-13b must be present. Transit timing variations can be used to constrain the mass and orbital elements of the perturbing planet, in our case those of HAT-P-13c.

The global modeling described in § 3.3 treats the transit centers of the FLWO 1.2 m telescope follow-up as independent variables, i.e., it automatically provides the basis for a TTV analysis. Throughout this work, by TTV we refer to the time difference between the observed transit center  $(T_{c,i})$  and the calculated value based on a fixed epoch and period as given in in Tab. 4, i.e. in the O-C sense. The resulting TTVs are shown in Fig. 7. We believe the error-bars to be realistic as they are the result of a full MCMC analysis, where all parameters are varied (including the EPD and TFA parameters). As further support for this, the transits around  $N_{tr} = 0$  $(T_{c,-1},T_{c,0},T_{c,1})$  show a standard deviation that is comparable to the error-bars (and we can safely assume zero TTV within a  $\pm 2.9163$  day time range). It is also apparent from the plot that the smallest error-bars correspond to the full transit observations  $(N_{tr} = 0 \text{ and } 24)$ . TTVs of the order of 100 seconds are seen from the best fit period. Given the large error-bars on our transit centers (of the order of 100 seconds), in part due to possible remaining systematics in the partial transit light curve events. we consider these departures suggestive only.

Nevertheless, it is tempting to compare our results with analytic formulae presented by Agol et al. (2005) and Borkovits et al. (2003). The HAT-P-13(b,c) system corresponds to case "ii" of Agol et al. (2005), i.e. with an exterior planet on an eccentric orbit having a much larger semi-major axis than the inner planet. Formulae for the general (non co-planar) case are given by Borkovits et al. (2003, Eq. 46). The TTV effect will depend on the following known parameters for the HAT-P-13 system:  $M_{\star}, M_{p}, P, i_{p}, P_{2}, e_{2}, \omega_{2}, m_{2} \sin i_{2}$  and  $T_{2,peri}$  (these are listed in Tab. 4 and Tab. 5). The TTV will also depend on the following unknown parameters: the true inclination of HAT-P-13c,  $i_2$ , and the relative angle of the orbital normals projected onto the plane of the sky,  $D = \Omega_1 - \Omega_2$  (see Fig. 1 of Borkovits et al. 2003). In addition to the gravitational perturbation of HAT-P-13c on HAT-P-13b, the barycenter of the inner subsystem (composed of the host star HAT-P-13, and the inner planet HAT-P-13b) orbits about the three-body barycenter due to the massive HAT-P-13c companion. This leads to light-travel time effects in the transit times of HAT-P-13b (TTVl) that are of the same order of magnitude as the TTV effect due to the perturbation (TTVg).

We have evaluated the analytic formulae including both the TTVg and TTVl effects for cases with  $i_2 =$  $83.4 \pm 0.6^{\circ}$  (corresponding to co-planar inner and outer orbits, and to  $M_2 = 15.2 M_J$ ), and  $i_2 = 8.1^{\circ}$  (an ad hoc value yielding an almost face-on orbit with  $M_2$  = 105  $M_{\rm J}$ ), and  $D = 0^{\circ}$  or  $D = 90^{\circ}$ . These four analytic models are illustrated in Fig. 7. The bottom panel of Fig. 7 shows the TTVl and TTVg effects separately for the  $i_2 = 83.4 \pm 0.6^{\circ}$  and  $D = 0^{\circ}$  case. The  $i_2 = 83.4 \pm 0.6^{\circ}$ ( $M_2 = 15.2 M_J$ ) cases give TTV variations of the order of 15 seconds. The functional form of the  $i_2 = 83.4 \pm 0.6^{\circ}$ and  $D = 0^{\circ}$  case appears to follow the observational values, albeit with much smaller amplitude. Curiously, for this case the TTVl effect cancels the TTVg effect to some extent (bottom panel of Fig. 7). Increasing the mass of the outer companion by decreasing  $i_2$  at constant  $m_2 \sin i_2$  does increase the TTV amplitude up to 100 seconds at  $i_2 = 8.1^{\circ}$ , but the functional form changes and no longer resembles the trend seen in the observational data.

In conclusion, while it is premature to fit the current data-set with analytic models because of the small number of data-points (7) and the large error-bars ( $\sim$ 100 sec), these data are not inconsistent with the pres-

TABLE 6 Transit timing variations of HAT-P-13.

$N_{tr}$	$\binom{T_c}{(\text{BJD})}$	$\sigma_{ m T_c} ( m sec)$	O-C (sec)
$-68 \\ -1 \\ 0 \\ 1 \\ 24 \\ 35 \\ 62$	$\begin{array}{c} 2454581.62406\\ 2454777.01287\\ 2454779.92953\\ 2454782.84357\\ 2454849.92062\\ 2454882.00041\\ 2454960.73968 \end{array}$	$105.7 \\86.7 \\54.3 \\133.6 \\65.1 \\129.5 \\154.1$	$\begin{array}{r} -10.9 \\ -58.5 \\ -24.2 \\ -215.6 \\ 50.7 \\ 132.2 \\ 156.1 \end{array}$

ence of TTVs, and will prove useful in later analyses. Future measurements of full transit events with high accuracy in principle can determine both  $i_2$  and D, i.e., HAT-P-13c may become an RV-based detection with known orbital orientation and a true mass (even if it does not transit).



FIG. 7.— (Top) Transit times of the individual transit events of HAT-P-13b. The filled circles correspond to the global analysis described in § 4.2. The large filled square, the open circle and the open square correspond to the apastron, conjunction and periastron of the perturber HAT-P-13c. Overlaid are analytic models for the transit timing variation due to gravitational effects (TTVg) and light-travel time effects (TTVl) for four scenarios: i) the inclination of HAT-P-13c is  $i_2 = 83.1^{\circ} = i_1$ , ii)  $i_2 = 8.1^{\circ}$  (at fixed  $m_2 \sin i_2$ , corresponding to a large mass for HAT-P-13c), iii)  $i_2 = 83.1^{\circ}$  and the mutual inclination of the orbital normals in the plane of the sky is  $D = \Omega_1 - \Omega_2 = 90^{\circ}$  (see Fig. 1 of Borkovits et al. (2003)), or iv)  $i_2 = 8.1^{\circ}$  and  $D = 90^{\circ}$ . (Bottom) The selected case "i" from above with zoomed-in vertical scale, showing the TTVg (dashed line) and TTVl (dotted line) effects separately, and their net effect (solid line).

#### 4.3. The planet HAT-P-13c

The probability of HAT-P-13c transiting the host star, as seen from the Earth, depends on  $R_{\star}$ ,  $R_{2p}$ ,  $e_2$ ,  $\omega_2$ , and  $a_2$  (Kane & von Braun 2009). We evaluated the transit probability in a Monte Carlo fashion, as part of the global modeling, resulting in  $P_{2,tr} = 0.0130 \pm 0.0008$ if  $R_{2,p} = 1 R_{\rm J} (P_{2,tr} = 0.0122 \pm 0.0007$  if  $R_{2,p} \rightarrow 0$ ). This derivation assumes an isotropic inclination distribution. However, dynamical constraints, such as precise measurements of TTV effects, or analysis of orbital stability, may further limit the allowed inclination range, and thus increase or decrease the chance of transits.

Unfortunately, our HATNet and FLWO 1.2 m datasets does not cover any time interval around the expected transit times of HAT-P-13c, thus we can not prove or refute the existence of such transits. Nevertheless, it is an interesting thought experiment to characterize the putative transits of HAT-P-13c, should they occur. HAT-P-13c would be the longest period transiting planet discovered to date, with a semi-major axis of over 1 AU, and a period of  $\sim 428 \,\mathrm{days}$ , about 4 times longer than the current record holder HD 80606 (Naef et al. 2001; Fossey, Waldmann, & Kipping 2009; Moutou et al. 2009). With a true mass of  $15.2 M_{\rm J}$ , we have good reasons to believe that its radius would be around  $1R_{\rm J}$ , based on heavy-mass transiting planets like HAT-P-2b  $(8.84 M_{\rm J})$ , Bakos et al. 2007), XO-3b  $(11.79 M_J, Johns-Krull et al.$ 2008), Corot-3b (21.66  $M_{\rm J}$ , Deleuil et al. 2008), all having radii around  $1R_{\rm J}$ . If the transits are full (i.e. not grazing), then the transit depth would be similar to that of the inner planet HAT-P-13b. The duration of the transit could be up to 30.8 hours. Follow-up observations of such transits would require either a world-wide effort, or uninterrupted space-based observations. The next transit center, according to the present analysis, will occur at 2009 April 12 9am UTC. Since we are looking at the orbit of HAT-P-13c along the semi-latus rectum (parallel to the minor-axis), the chance for an occultation of the planet by the star has a very similar chance of occurance as the primary eclipse. Observations of the secondary eclipse could greatly decrease the error on  $e_2$  and  $\omega_2$ .

- Agol, E., Steffen, J., Sari, R., & Clarkson, W. 2005, MNRAS, 359, 567
- Cox, A. N. 2000, Allen's Astrophysical Quantities
- Bakos, G. Á., Lázár, J., Papp, I., Sári, P. & Green, E. M. 2002, PASP, 114, 974
- Bakos, G. Á., Noyes, R. W., Kovács, G., Stanek, K. Z., Sasselov, D. D., & Domsa, I. 2004, PASP, 116, 266
- Bakos, G. Á., et al. 2007, ApJ, 670, 826
- Bakos, G. Á., et al. 2009, arXiv:0901.0282
- Baraffe, I., Chabrier, G., & Barman, T. 2008, A&A, 482, 315
- Barning, F. J. M. 1963, Bull. Astron. Inst. Netherlands, 17, 22
- Bodenheimer, P., Lin, D. N. C., & Mardling, R. A. 2001, ApJ, 548, 466
- Borkovits, T., Érdi, B., Forgács-Dajka, E., & Kovács, T. 2003, A&A, 398, 1091
- Burke, C. J., et al. 2007, ApJ, 671, 2115
- Butler, R. P. et al. 1996, PASP, 108, 500
- Carpenter, J. M. 2001, AJ, 121, 2851
- Carter, J. A., Yee, J. C., Eastman, J., Gaudi, B. S., & Winn, J. N. 2008, ApJ, 689, 499
- Claret, A. 2004, A&A, 428, 1001
- Deleuil, M., Deeg, H. J., Alonso, R., Bouchy, F., & Rouan, D. 2008, arXiv:0810.0919
- Deeming, T. J. 1975, Ap&SS, 36, 137
   Droege, T. F., Richmond, M. W., Sallman, M. P., & Creager, R. P. 2006, PASP, 118, 1666
- Fabrycky, D. C., Johnson, E. T., & Goodman, J. 2007, ApJ, 665, 754
- Fabrycky, D. C. 2009, IAU Symposium, 253, 173
- Ford, E. 2006, ApJ, 642, 505
- Fortney, J. J., Marley, M. S., & Barnes, J. W. 2007, ApJ, 659, 1661
- Fossey, S. J., Waldmann, I. P., & Kipping, D. M. 2009, MNRAS, 396, L16

As regards to the nature of HAT-P-13c, even at its minimal mass  $(15.2 \pm 1.0 M_{\rm J})$  it is the 10th most massive planet out of 327 planets listed on the Extrasolar Planet Encyclopedia as of 2009 July. Curiously, the recently announced Doppler-detection HD 126614b (Howard et al. 2009) appears to have similar orbital characteristics to HAT-P-13c. Both are Jovian planets in P > 1 yr, e = 0.7 orbits around metal-rich ([Fe/H]  $\approx 0.5$ ) stars. As described earlier in  $\S$  4.2, we have good hopes that in the near future precise TTV measurements of the inner planet HAT-P-13b, will constrain the orbital inclination and thus the true mass of HAT-P-13c. Further, such TTV variations can also constrain the mutual inclination of HAT-P-13b and HAT-P-13c. Measuring the sky-projected angle between the stellar spin axis and the orbital normal of HAT-P-13b (the inner planet) via the Rossiter-McLaughlin effect will shed light on the migration history of HAT-P-13b, and by inference the scattering history between HAT-P-13b and HAT-P-13c.

HATNet operations have been funded by NASA grants NNG04GN74G, NNX08AF23G and SAO IR&D grants. Work of G.A.B. and J. Johnson were supported by the Postdoctoral Fellowship of the NSF Astronomy and Astrophysics Program (AST-0702843 and AST-0702821, respectively). We acknowledge partial support also from the Kepler Mission under NASA Cooperative Agreement NCC2-1390 (D.W.L., PI). G.K. thanks the Hungarian Scientific Research Foundation (OTKA) for support through grant K-60750. This research has made use of Keck telescope time granted through NOAO (program A146Hr, A264Hr) and NASA (N128Hr, N145Hr).

## REFERENCES

- Hansen, B. M. S., & Barman, T. 2007, ApJ, 671, 861
- Hartman, J. D., et al. 2009, arXiv:0904.4704
- Holman, M. J., & Murray, N. W. 2005, Science, 307, 1288
- Holman, M. J., et al. 2006, ApJ, 652, 1715
- Howard, W. A., et al. 2009, submitted to ApJ
- Kane, S. R., & von Braun, K. 2009, IAU Symposium, 253, 358
- Kovács, G., Zucker, S., & Mazeh, T. 2002, A&A, 391, 369
- Kovács, G., Bakos, G. Á., & Noyes, R. W. 2005, MNRAS, 356, 557 Johns-Krull, C. M., et al. 2008, ApJ, 677, 657
- Latham, D. W. 1992, in IAU Coll. 135, Complementary Approaches to Double and Multiple Star Research, ASP Conf. Ser. 32, eds. H. A. McAlister & W. I. Hartkopf (San Francisco: ASP), 110
- McCullough, P. R., et al. 2006, ApJ, 648, 1228
- Pál, A., & Bakos, G. Á. 2006, PASP, 118, 1474
- Pál, A., Bakos, G. Á., Noyes, R. W. & Torres, G. 2008, proceedings of IAU Symp. 253 "Transiting Planets", ed. by F. Pont (arXiv:0807.1530)
- Pál, A. 2008b, MNRAS, 390, 281
- Pál, A., PhD thesis, (arXiv:0906.3486)
- Press, W. H., Teukolsky, S. A., Vetterling, W. T. & Flannery, B. P., 1992, Numerical Recipes in C: the art of scientific computing, Second Edition, Cambridge University Press
- Mandel, K., & Agol, E. 2002, ApJ, 580, L171
- Mardling, R. A. 2007, MNRAS, 382, 1768
- Marcy, G. W., & Butler, R. P. 1992, PASP, 104, 270
- McLaughlin, D. B. 1924, ApJ, 60, 22
- Miller, N., Fortney, J. J., & Jackson, B. 2009, arXiv:0907.1268
- Moutou, C., et al. 2009, A&A, 498, L5
- Naef, D., et al. 2001, A&A, 375, L27
- Shporer, A., et al. 2008, arXiv:0806.4008
- Skrutskie, M. F., et al. 2006, AJ, 131, 1163
- Smith, A. M. S., et al. 2009, arXiv:0906.3414

- Sozzetti, A. et al. 2007, ApJ, 664, 1190
  Torres, G., Boden, A. F., Latham, D. W., Pan, M. & Stefanik, R. P. 2002, AJ, 124, 1716
  Torres, G. et al. 2007, ApJ, 666, 121
  Valenti, J. A., & Fischer, D. A. 2005, ApJS, 159, 141
  Valenti, J. A., & Piskunov, N. 1996, A&AS, 118, 595

- Vogt, S. S. et al. 1994, Proc. SPIE, 2198, 362 Yi, S. K. et al. 2001, ApJS, 136, 417 Winn, J. N., et al. 2007, AJ, 133, 1828 Wright, J. T., et al. 2007, ApJ, 657, 533